Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

there are faint lines reaching from the A. M. E. down upon the mandibles, sternum reddish brown on the sides, paler in the center; legs reddish, more yellow at tips, a few dark spots at the base of hairs, dorsum of abdomen reddish brown, often with a light median stripe enclosing a spear-mark at base, and a short light mark on each side near tip; venter yellowish with a broad median brown stripe, spinnerets brown. The ♂ is darker than the ♀. Sometimes the legs are a little banded. Cephalothorax highest at eye-region, gradually sloping concavely until near the posterior end, where if suddenly curves down; legs spiny, abdomen widest in front, tapering to apex; the epigynum consists of a rounded finger, somewhat similar to *O. cinerea* but more slender. Male palpal organ black, the tarsus is more slender than in the other species, the tibia has on the inner side a short pointed projection similar to *O. salticus*, the basal part of the palpal organ is more complicated than in that species.

The four species of *Oxyopes* may be separated by the following table:

| | | |
|---|---|---|
| 1 | Femora with a black line on under-side . . . . . | **salticus** |
| | Femora without a black line on under-side . . . . . | 2 |
| 2 | Abdomen light, with black side and median stripes . . | **scalaris** |
| | Abdomen dark, with a few light spots . . . . . | 3 |
| 3 | Dorsum of cephalothorax straight, about as high in middle as in eye-region | **cinerea** |
| | Dorsum of cephalothorax concave, higher in eye-region than in middle | **rufipes** |

## LOCAL ENTOMOLOGICAL NOTES.

Members of the New York Entomological Society and all others, are solicited to contribute to this column, their rare captures, local lists and other items of interest relating to the insect fauna of New York city and vicinity.

## LIST OF THE COLEOPTERA OF NORTH EASTERN AMERICA,

### WITH SPECIAL REFERENCE TO THE FAUNA OF NEW YORK CITY AND VICINITY.

By Charles W. Leng and Wm. Beutenmuller.

(Continued from page 96.)

Schizogenius *Putz.*

**S. planulatus** *Lec.*—N. Y. Taken by Mr. Linell at Coney Island.

**S. lineolatus** *Say.*—N. E. Amer. Lives under stones along river banks, June and Sept. Not common in this vicinity.

**S. ferrugineus** *Putz.*—N. E. Amer. Occurs in salt marshes. Not common in this vicinity.

**S. amphibius** *Hald.*—N. Y., Mo. Taken in this vicinity by Mr. Wm. Jülich.

## ARDISTOMIS *Putz.*

**A. obliquata** *Putz.*—N. J. and southward. Taken at Atlantic City, N. J., by Mr. Liebeck.

**A. morio** *Dej.*—N. Y. to Fla. Taken in this vicinity by Mr. Jülich.

**A. viridis** *Say.*—N. E. Amer. Found in sandy banks. Not common in this vicinity.

**A. puncticollis** *Putz.*—Ohio (Dury).

## PANAGÆUS *Lat.*

**P. crucigerus** *Say.*—N. Y., N. J. and Southward. Rare in this vicinity. May und June under stones. Taken in April in deserted ant hills (W. B).

**P. fasciatus** *Say.*—N. Y., N. J. and southward. Very rare in this vicinity. May and June.

## NOMIUS *Lap.*

**N. pygmæus** *Dej.*—Taken at Newark, N. J., by Mr. Bischoff. Also found in Canada (Harrington).

## BEMBIDIUM *Lat.*

**B. punctatostriatum** *Say.*—N. Y., N. J., O. Found along the banks of the Passaic, N. J., in May and Sept.

**B. impressum** *Fabr.*—Buffalo, N. Y.

**B. paludosum** *Sturm.*—Buffalo, N. Y.

**B. inæquale** *Say.*—N. E. Amer. Found along river banks in May and Sept.

**B. nitidulum** *Dej.*—N. E. Amer. Found along the banks of the Passaic, N. J., in May and Sept.

**B. nitidum** *Kby.*—Buffalo, N. Y.

**B. americanum** *Dej.*—N. E. Amer. Rather common in this vicinity, along banks of ponds. May and Sept.

**B. antiquum** *Dej.*—N. Y., N. J. Along river banks, not common in this vicinity.

**B. chalceum** *Dej.*—N. Y., N. J., Ohio. Found along the banks of the Passaic in May and Sept.

**B. concolor** *Kby.*—Buffalo, N. Y.

**B. nigrum** *Say.*—N. Y., N. J., Can. Found along the banks of the Passaic, N. J., in May and Sept.

**B. planatum** *Lec.*—N. J. Taken in the same locality as the preceding species.

**B. Kuprianovi** *Mann.*—Buffalo, N. Y.

**B. fugax** *Lec.*—Buffalo, N. Y., Mo.

**B. planum** *Hald.*—Canada, Buffalo, N. Y.

**B. rupestre** *Dej.*—N. E. Amer. Common in this vicinity.

**B. plagiatum** *Zimm.*—Buffalo, N. Y.

**B. lacunarium** *Zimm.*—N. Y., N. J. Found along brooks at Fort Lee, N. J. (Linell) and at Coney Island (Leng).

**B. dorsale** *Say.*—Can., N. Y., N. J., Ohio.

**B. patruele** *Dej.*—N. Y., N. J., Ohio, Pa. Found along river banks, May and Sept.

**B. variegatum** *Say.*—N. E. Amer. Common in this vicinity.

*var.* **posticum** *Lec.*—Taken at Coney Island, N. Y. (Leng).

**B. intermedium** *Kby.*—N. Y. (Buffalo) N. J., Ohio, Canada.

**B. nigripes** *Kby.*—Pa., (also Or. Br. Col).

**B. versicolor** *Lec.*—N. Y., N. J., Ohio. Found along the banks of the Passaic, N. J., in May and Sept.

**B. constrictum** *Lec.*—N. Y., N. J. Taken at Coney Island (Leng). Common in N. J. Sept.

**B. contractum** *Say.*—N. Y., N. J. Found along the seashore.

**B. affine** *Say.*—N. Y., N. J., Ohio.

**B. assimile** *Gyll.*—Buffalo, N. Y.

**B. quadrimaculatum** *Linn.*—N. E. Amer. Common everywhere, in fields and gardens.

**B. semistriatum** *Hald.*—N. Y., N. J. May and Sept.

**B. picipes** *Kby.*—Ohio (Dury), Peekskill, N. Y. (Sherman), N. Hampshire.

**B. cordatum** *Lec.*—Ohio (Dury), also Mo.

**B. lævigatum** *Lec.*—Ohio, (Dury), also Mo.

**B. bifossulatum** *Lec.*—Peekskill, N. Y., (Sherman).

**B. bimaculatum** *Kby.*—Ottawa, Canada (Harrington).

**B. pedicellatum** *Lec.*—Pa. (also Missouri).

## TACHYS *Schaum.*

**T. proximus** *Say.*—Ohio, Buffalo, N. Y., N. J., Pa. Found along the banks of the Passaic, N. J. May and Sept.

**T. scitulus** *Lec.*—Ohio, Buffalo, N. Y., N. J. Found in the same locality as the preceding.

**T. occultator** *Casey.*—N. J. Taken at Brigantine Beach by Dr. Hamilton in September.

**T. lævus** *Say.*—Ohio, Buffalo, N. Y., N. J., Pa. Rare in this vicinity.

**T. nanus** *Gyll.*—N. E. Amer. Common in this vicinity under bark of trees; runs very fast.

**T. flavicauda** *Say.*—N. E. Amer. Common under bark of trees. May and September.

**T. tripunctatus** *Say.*—N. Y., N. J., Pa. Taken along the banks of the Passaic, N. J., in May and Sept.

**T. vivax** *Lec.*—Ohio, Buffalo, N. Y., Pa.

**T. capax** *Lec.*—N. J. and westward. Found in cranberry bogs in N. J. (Smith).

**T. xanthopus** *Say.*—N. Y., N. J. and westward.

**T. incurvus** *Say.*—N. E. Amer. Common in the vicinity of N. Y. City.

**T. fuscicornis** *Chd.*—N. J.

**T. pulchellus** *Lec.*—Ohio and westward.

**T. ventricosus** *Lec.*—Peekskill, N. Y. (Sherman).

## PATROBUS *Dej.*

**P. longicornis** *Say.*—N. E. Amer. Found in woods in moist places.

**P. septentrionis** *Dej.*—Northern States and Europe. Not found in this vicinity.

**P. rugicollis** *Rand.*—N. Y., Pa., Mass. and northward. Not found in this vicinity.

## PERICOMPSUS *Lec.*

**P. ephippiatus** *Say.*—N. Y. and westward. Found in sandy places near the Hudson River, N. Y. (Sherman).

## POGONUS *Dej.*

**P. texanus** *Chd.*—N. J. Rare at Atlantic City.

## TRECHUS *Clair.*

**T. rubens** *Fabr.*—Nova Scotia, N. Europe.

**T. chalybæus** *Mann.*—Alaska to N. Hampshire.

## MYAS *Dej.*

**M. cyanescens** *Dej.*—N. Y., N. J. Taken in this vicinity at Fort Lee and Clifton, N. J., and on Staten Island, in June and July. Rare.

**M. coracinus** *Say.*—N. Y., N. J., Ohio. Rare.

## PTEROSTICHUS *Bon.*

**P. adoxus** *Say.*—N. E. Amer. Not common in this vicinity.

**P. rostratus** *Newm.*—Can., N. Y. and southward.

**P. diligendus** *Chaud.*—N. E. Amer. Not common in this vicinity.

**P. honestus** *Say.*—N. E. Amet. Not rare in this vicinity.

**P. lachrymosus** *Newm.*—N. Y. and southward. Not rare in this vicinity.

**P. stygicus** *Say.*—N. E. Amer. Abundant in this vicinity.

**P. coracinus** *Newm.*—N. E. Amer. Not rare at Snake Hill, and Short Hills, N. J.

**P. relictus** *Newm.*—N. Y., Pa., Mich. Rare.

**P. superciliosus** *Say.*—N. Y., Pa., Mich. Rare.

**P. moestus** *Say.*—N. Y., N. J., Pa., W. Va. Rare in this vicinity.

**P. punctatissimus** *Rand.*—Mass. and Westward.

**P. sculptus** *Lec.*—N. Y. and southward. Taken at Atlantic City, N. J. (Smith).

**P. Hamiltoni** *Horn.*—Pa., E. Ohio, Md., W. Va. Not common.

**P. Sayi** *Brulle.*—N. E. Amer. Common in this vicinity.

**P. lucublandus** *Say.*—N. E. Amer. Very common everywhere.

**P. ebeninus** *Dej.*—N. Y. and southward. Rare in this vicinity.

**P. caudicalis** *Say.*—N. E. Amer. Rare in this vicinity.

**P. luctuosus** *Dej.*—N. E. Amer. Taken at Snake Hill, N. J. Rare.

**P. corvinus** *Dej.*—N. E. Amer. Not common in this vicinity.

**P. purpuratus** *Lec.*—N. Y. and westward. Not found in this vicinity. Rare.

**P. mutus** *Say.*—N. E. Amer. Very common everywhere in this vicinity.
**P. pennsylvanicus** *Lec.*—N. E. Amer. Not found in this vicinity.
**P. orinomum** *Leach.*—Canada, Nova Scotia.
**P. Luczotii** *Dej.*—N. Y. and northward. Not found in this vicinity.
**P. erythropus** *Dej.*—N. E. Amer. Rare in this vicinity.
**P. patruelis** *Dej.*—N. E. Amer. Common in this vicinity.
**P. femoralis** *Kby.*—Can., N. Y. and westward. Wis. (O. Dietz).
**P. corrusculus** *Lec.*—N. Y., Mass. Rare under moss.
**P. mandibularis** *Kby.*—N. Hampshire, Can.
**P. Haldemani** *Lec.*—N. Y. and southward. Rare on Staten Island (Leng).
**P. tartaricus** *Say.*—N. Y., southward and westward. Taken at Hoboken, N. J. and Astoria, L. I. May.
**P. scrutator** *Lec.*—N. Y., Can., Ill.
**P. gravis,** *Lec.*—Pa.
**P. permundus** *Say.*—Ohio.
**P. substriatus** *Lec.*—N. J. (Davis), N. Y. (Jülich) and southward.
**P. rotundatus** *Lec.*—N. Y. (Jülich), Ga. (Lec).
**P. unicolor** *Say.*—N. Y. and southward. Not found in this vicinity.
**P. approximatus** *Lec.*—Pa. and southward.
**P. hudsonicus** *Lec.*—Labrador.

## Evarthrus *Lec.*

**E. sigillatus** *Say.*—N. J. and southward.
**E. sodalis** *Lec.*—Pa. and southward. Wis., Ohio (Dietz).
**E. furtivus** *Lec.*—Pa. and southward.

## Amara *Bon.*

**A. avida** *Say.*—N. E. Amer. Rare in this vicinity.
**A. arenaria** *Lec.*—Buffalo, N. Y., N. Hampshire.
**A. exarata** *Dej.*—N. E. Amer. Not common.
**A. latior** *Kby.*—N. J.
**A. angustata** *Say.*—N. E. Amer. Not rare in this vicinity.
**A. pallipes** *Kby.*—N. J. and westward.
**A. impuncticollis** *Say.* N. E. Amer. Common in this vicinity.
**A. basillaris** *Say.*—N. Y., N. J. (Linell).
**A. cupreolata** *Putz.*—N. J., Ill.
**A. interstitialis** *Dej.*—Can., N. Y., N. J. Not rare in this vicinity.
**A. obesa** *Say.*—N. Y., N. J., Can. and westward.
**A. fallax** *Lec.*—Buffalo, N. Y. and westward.
**A. erratica** *Sturm.*—Can., Vt., Mich.
**A. chalcea** *Dej.*—N. Y., N. J. and westward.
**A. gibba** *Lec.*—N. J.
**A. rubrica** *Hald.*—N. Y., N. J. Rare in this vicinity.
**A. subænea** *Lec.*—Can., N. Y., N. J.
**A. musculus** *Say.*—N. Y., N. J. and westward. Not common in this vicinity.
**A. acutangula** *Putz.*—N. J. Rare.
**A. similis** *Kby.*—Labrador.

**A. polita** *Lec.*—Can., N. Y. (O. Dietz).
**A. septentrionalis** *Lec.*—N. Y. (O. Dietz).
**A. hyperborea** *Dej.*—Labrador.

## LOXANDRUS *Lec.*

**L. agilis** *Dej.*—Pa. and southward.

## DIPLOCHILA *Brulle.*

**D. laticollis** *Lec.*—New York and westward. Not rare in moist places, in this vicinity. May and June.
*var.* **major** *Lec.*—Found with the preceding.
**D. impressicollis** *Dej.*—Can., N. Y. and westward.

## DICÆLUS *Bon.*

**D. dilatatus** *Say.*—N. Y. and westward. Rare in this vicinity. May and June.
**D. purpuratus** *Bon.*—N. Y., Mass., southward and westward.
**D. sculptilis** *Say.*—Ohio, southward and westward. Wis. (O. Dietz).
**D. furvus** *Dej.*—Ohio, Southward and westward.
**D. crenatus** *Lec.*—A southern species on the Buffalo list.
**D. ovalis** *Lec.*—N. Y., west and southward. Taken at Snake Hill, N. J.
**D. elongatus** *Bon.*—N. E. Amer. Not rare in this vicinity.
**D. ambiguus** *Laf.*—N. Y., west and south. Not found in this vicinity.
**D. teter** *Bon.*—N. E. Amer. Not rare along the Palisades, N. J.
**D. politus** *Dej.*—N. E. Amer. Not rare in this vicinity.

## LICINUS *Lat.*

**L. silphoides** *Fab.*—Mass. and Europe.

## BADISTER *Clairv.*

**B. notatus** *Hald.*—Can., N. Y., N. J., westward and southward. Rare in this vicinity.
**B. micans** *Lec.*—Mass., N. Y., N. J., Ill., Fla. Rare in this vicinity.
**B. pulchellus** *Lec.*—Can., N. Y., Ill., Ind., Fla.
**B. maculatus** *Lec.*—Pa.
**B. ferrugineus** *Dej.*—Cal., Alaska (on the Buffalo list).
**B. flavipes** *Lec.*—N. Y., La., Fla.
**B. reflexus** *Lec.*—N. Y., Mich., La.

## CALATHUS *Bon.*

**C. gregarius** *Say.*—Can., N. Y. to Fla. to Tex. Common in various localities in this vicinity.
**C. ingratus** *Dej.*—Can., Labrador, Alaska.
**C. opaculus** *Lec.*—N. E. Amer. Rare in this vicinity.
**C. advena** *Lec.*—Alaska to N. Hampshire.
**C. impunctatus** *Say.*—N. E. Amer. Not rare in this vicinity, especially at the sea shore.

## PLATYNUS *Bon.*

**P. angustatus** *Dej.*—Middle and Southern States. Rare at Fort Lee, N. J.

**P. caudatus** *Lec.*—Ohio.

**P. hypolithus** *Say.*—Middle and Southern States.

**P. decens** *Say.*—Can. to N. C. Scarce in this vicinity.

**P. sinuatus** *Dej.*—N. E. Amer. Not common in this vicinity.

**P. opaculus** *Lec.*—Middle and Western States. Rare at Fort Lee, N. J.

**P. tenuicollis** *Lec.*—Lake Superior. Taken at Atlantic City, N. J: (Castle).

**P. cincticollis** *Say.*—Middle and Southern States. Common under stones along brooks.

**P. reflexus** *Lec.*—Can. and Middle States. Found in same situation as the preceding species.

**P. extensicollis** *Say.*—N. E. Amer. Common under stones along brooks and rivers in this vicinity.

*var.* **viridis** *Lec.*—Southern and Western States. (On the Buffalo list).

**P. decorus** *Say.*—N. E. Amer. Not rare along the Passaic, N. J.

**P. anchomenoides** *Rand.*—Can. and Maine.

**P. clemens** *Lec.*—Nova Scotia.

**P. errans** *Say.*—Canada.

**P. moerens** *Dej.*—Can. and Atlantic States.

**P. pusillus** *Lec.*—Can., N. Y. and westward.

**P. collaris** *Say*—Southern States. (On the Buffalo list).

**P. tenuis** *Lec.*—Can. and Middle States.

**P. atratus** *Lec.*—N. E. Amer. Rare in this vicinity.

**P. melanarius** *Dej.*—N. E. Amer. Common in this vicinity.

**P. propinquus** *G. & H.*—Can. to Mass., N. J.

**P. affinis** *Kby.* N. E. Amer. Rare in this vicinity.

**P. corvus** *Lec.*—Can. to Or.

**P. metallescens** *Lec.*—Lake Superior to N. J. Not rare in this vicinity.

**P. deceptivus** *Lec.*—Nova Scotia to Lake Superior.

**P. cupripennis** *Say.*—N. E. Amer. Common in this vicinity.

*var.* **nitidulum** *Dej.*—Bear Lake, Pa.

**P. Hardyi** *Lec.*—Newfoundland.

**P. excavatus** *Dej.*—Can., Middle and Western States. Rare in this vicinity.

**P. ferreus** *Hald.*—Middle and Western States. Not rare along the Palisades, N. J.

**P. basalis** *Lec.*—Pa., N. J., to Fla. Very rare. Taken at Hoboken, N. J. (Linell).

**P. maculicollis** *Dej.*—Cal. (On the Buffalo list).

**P. variolatus** *Lec.*—Cal. (On the Buffalo list).

**P. nutans** *Say.*—Can., N. Y. to Fla. Rare in this vicinity.

**P. octopunctatus** *Fabr.*—N. E. Amer. Rare in this vicinity.

**P. placidus** *Say.*—N. E. Amer. Common.

**P. vicinus** *G. & H.*—Can., Lake Superior.

**P. obsoletus** *Say.*—N. E. Amer. Rather common in this vicinity.

**P. quadripunctatus** *Dej.*—Can., Alaska, N. Y. (On the Buffalo list).

**P. æruginosus** *Dej.*—Atlantic States. Not common under bark in this vicinity.

**P. crenistriatus** *Lec.*—N. Y. and westward. Abundant on seashore of Long Island and New Jersey.

**P. rubripes** *Zimm.*—Middle and Western States. Not common in this vicinity.

**P. punctiformis** *Say.*—N. E. Amer. Rather common in this vicinity.

**P. sordens** *Kby.*—Can., to N. J. Rare in this vicinity.

**P. picicornis** *Lec.*—Can., N. Y. and westward. Not found around N. Y. City.

**P. ruficornis** *Lec.*—N. Y. and Westward. Wis. (O. Dietz).

**P. retractus** *Lec.*—Can., Pa., Mass. Lake Superior.

**P. picipennis** *Kby.*—Mass., N. Y., N. J. and northward.

**P. lutulentus** *Lec.*—N. E. Amer. Not common in this vicinity.

**P. exaratus** *Mann.*—Pa.

**P. limbatus** *Say.*—Can. (Couper). Mo.

### Olisthopus *Dej.*

**O. parmatus** *Say.*—N. E. Amer.

**O. micans** *Lec.*—N. J. southward and westward.

### Atranus *Lec.*

**A. pubescens** *Dej.*—N. Y. and westward.

### Leptotrachelus *Lat.*

**L. dorsalis** *Fab.*—N. Y. and westward.

### Casnonia *Lat.*

**C. pennsylvanica** *L.*—N. E. Amer.

**C. ludoviciana** *Salle.*—D. C. and South.

### Galerita *Fab.*

**G. janus** *Fab.*—N. E. Amer. Common everywhere.

**G. bicolor** *Dru.*—N. Y., N. J., Ga., Mo.

### Tetragonoderus *Dej.*

**T. fasciatus** *Hald.*—N. Y. and westward.

### Lebia *Lat.*

**L. grandis** *Hentz.*—N. E. Amer. Common everywhere.

**L. atriventris** *Say.*—N. E. Amer. Not as common as the preceding species.

**L. tricolor** *Say.*—N. E. Amer.

**L. pulchella** *Dej.*—N. E. Amer. Not rare.

**L. cyanipennis** *Dej.*—N. J., Cal.

**L. marginicollis** *Dej.*—N. J. Rare.

**L. viridis** *Say.*—N. E. Amer. Common.

**L. moesta** *Lec.*—N. Y., N. J. and westward.

**L. pumila** *Lec.*—N. E. Amer. Not common.
**L. pleuritica** *Lec.*—N. E. Amer. Not common.
**L. viridipennis** *Dej.*—N. E. Amer.
**L. lobulata** *Lec.*—Ohio, Va., La.
**L. ornata** *Say.*—N. E. Amer.
**L. collaris** *Dej.*—Middle and Southern States.
**L. analis** *Dej.*—N. E. Amer.
**L. fuscata** *Dej.*—N. E. Amer.
*var.* **frigida** *Chaud.*—Mass.
**L. scapularis** *Dej.*—N. E. Amer.
**L. vittata** *Fab.*—N. Y., N. J., Pa. to Tex. Not common in this vicinity.
*var.* **Spraguei** *Horn.*—N. J. (Castle), Tex.
**L. furcata**—*Lec.* N. E. Amer.
**L. bivittata** *Fab.*—Mass., N. Y., N. J. Westward and Southward.

## COPTODERA *Dej.*

**C. aerata** *Dej.*—N. E. Amer. Rare.

## DROMIUS *Bon.*

**D. piceus** *Dej.*—Mass., N. Y., Pa. to Cal. Found under bark and on flowers.

## APRISTUS *Chd.*

**A. cordicollis** *Lec.*—N. Y., N. J.
**A. subsulcatus** *Dej.*—N. Y., Can.

## BLECHRUS *Mots.*

**B. nigrinus** *Mann.*—Can., N. Y. and westward. Found under bark.

## METABLETUS *Sch.-Goeb.*

**M. americanus** *Dej.*—N. E. Amer.

## AXINOPALPUS *Lec.*

**A. biplagiatus** *Dej.*—Can., Mass. to Calif.

## CALLIDA *Dej.*

**C. viridipennis** *Say.*—N. Y. to Texas.
**C. punctata** *Lec.*—N. J., N. Y., Mich., O., La., Ks., Can.
**C. purpurea** *Say.*—N. J., N. Y., Mich., Ga., Mo., Ks., Neb.

## PLOCHIONUS *Dej.*

**P. Bonfilsii** *Dej.*—Pa., Mex.
**P. timidus** *Hald.*—N. Y., N. J., Ala., Cal. Under bark.
**P. amandus** *Hald.*—N. Y. and Southward.

### PINACODERA *Schaum.*

**P. limbata** *Dej.*—N. Y., N. J.
*var.* **fuscata** *Dej.*—N. Y.
**P. platicollis** *Say.*—N. Y., N. J., Fla., Tex.

### CYMINDIS *Lat.*

**C. cribricollis** *Dej.*—N. Y. and westward.
**C. elegans** *Lec.*—N. Y.
**C. unicolor** *Kby.*—L. Sup., Labrador.
**C. americana** *Dej.*—N. Y., N. J. and westward.
**C. pilosa** *Lec.*—Can., N. Y., N. J. and westward.
**C. borealis** *Lec.*—Can., Nova Scotia.
**C. neglecta** *Hald.*—Can., N. Y., N. J., Pa.

### APENES *Lec.*

**A. lucidula** *Dej.*—N. Y., N. J., Fla. Seashore.
**A. sinuata** *Say.*—N. Y., N. J.

### HELLUOMORPHA *Lap.*

**H. bicolor** *Harr.*—Mass. to Fla. and westward. Taken at Mosholu, N. Y., and Fort Lee, N. J. Rare.
**H. ferruginea** *Lec.*—N. Y. to Tex. Rare in this vicinity.

### BRACHYNUS *Web.*

**B. viridipennis** *Dej.*—N. Y., N. J.
**B. minutus** *Harr.*—N. Y., N. J.
**B. perplexus** *Dej.*—Can., N. Y., N. J. and westward.
**B. medius** *Harr.*—N. E. Amer.
**B. quadripennis** *Dej.*—N. Y., N. J., Pa., Mass.
**B. conformis** *Dej.*—N. Y., N. J.
**B. cyanipennis** *Say.*—N. Y., N. J.
**B. alternans** *Dej.*—Can., N. Y., N. J. and westward.
**B. fumans** *Fab.*—N. E. Amer.
*var.* **similis** *Lec.*—N. E. Amer.
**B. cordicollis** *Dej.*—Can., N. Y., N. J. and westward.
**B. americanus** *Lec.*—Can., N. Y. and westward.
**B. ballistarius** *Lec.*—Can., Ill., Mo.

### CHLAENIUS *Bon.*

**C. erythropus** *Germ.*—Ohio to Neb., La. Rare in N. J., Buffalo, N Y., Fla. (O. Dietz).
**C. fuscicornis** *Dej.*—Gulf States to Ill. (On the Buffalo list.)
**C. sericeus** *Forst.*—N. E. Amer. Common everywhere.
**C. laticollis** *Say.*—N. Y. to Fla. and westward.
**C. diffinis** *Chd.*—Middle States Region.
**C. aestivus** *Say.*—N. E. Amer.
**C. prasinus** *Dej.*—Middle States to Col.
**C. leucoscelis** *Chev.*—N. E. Amer. Under stones by Croton River, N. Y.

**C. solitarius** *Say.*—N. E. Amer. Not found in this vicinity.
**C. vafer** *Lec.*—Texas. (On the Buffalo list.)
**C. nemoralis** *Say.*—N. E. Amer. Common.
**C. tricolor** *Dej.*—N. E. Amer. Common everywhere.
**C. pennsylvanicus** *Say.*—N. Y., N. J. and northward.
**C. impunctifrons** *Say.*—N. E. Amer. Not common.
**C. niger** *Rand.*—N. E. Amer. Rare.
**C. purpuricollis** *Rand.*—N. E. Amer. Very rare.
**C. tomentosus** *Say.*—N. E. Amer. Common.
**C. platyderus** *Chd.*—Ill., west and southward.

### ANOMOGLOSSUS *Chd.*

**A. emarginatus** *Say.*—N. E. Amer. Not rare in this vicinity.
**A. pusillus** *Say.*—N. E. Amer. Rare in this vicinity.

### BRACHYLOBUS *Chd.*

**B. lithophilus** *Say.*—N. E. Amer. Rare in this vicinity.

### LACHNOCREPIS *Lec.*

**L. parallelus** *Say.*—N. Y. and westward.

### OODES *Bon.*

**O. amaroides** *Dej.*—N. Y., N. J., Pa., southward and westward. Very rare in this vicinity.
**O. americanus** *Dej.*—N. Y. and southwasd.
**O. fluvialis** *Lec.*—Can., N. Y., southward and westward.
**O. Lecontei** *Chd.*—La., (On the N. J. list.)
**O. cupraeus** *Chd.*—Tex., Mo. (On the Buffalo list.)

### GEOPINUS *Lec.*

**G. incrassatus** *Dej.*—N. Y., N. J. and westward. Sandy districts near water, often six inches deep.

### NOTHOPUS *Lec.*

**N. zabroides** *Lec.*—Ill. (O. Dietz.)

### CRATACANTHUS *Dej.*

**C. dubius** *Dej.*—N. Y., N. J. and westward. Found in sandy places, near water.

### AGONODERUS *Dej.*

**A. lineola** *Fab.*—N. E. Amer. Common everywhere.
**A. infuscatus** *Dej.*—N. Y. and southward.
**A. pallipes** *Fab.*—N. E. Amer. Common everywhere.
**A. partiarius** *Say.*—N. E. Amer.
**A. pauperculus** *Dej.*—Southern States. (On the N. J. and Buffalo lists).
**A. indistinctus** *Dej.*—N. Y., N. J.
**A. testaceus** *Dej.*—N. Y., N. J.
**A. micros** *Lec.*—Eastern States.

## Discoderus *Lec.*

**D. parallelus** *Hald.*—N. J., southward and westward. Seashore.
**D. impotens** *Lec.*—Ohio. (O. Dietz.)

## Gynandropus *Dej.*

**G. hylacis** *Say.*—N. Y., N. J. Under bark.

## Harpalus *Lat.*

**H. dichrous** *Say.*—N. Y., N. J.
**H. vulpeculus** *Say.*—N. Y., N. J.
**H. autumnalis** *Say.*—N. Y., N. J. Rare at seashore.
**H. erraticus** *Say.*—Can., N. H., Mass., N. Y., N. J.
**H. viridiæneus** *Beauv.*—N. E. Amer.
**H. basilaris** *Kby.*—Can.
**H. laticeps** *Lec.*—Wis. (O. Dietz.) Can.
**H. caliginosus** *Fab.*—N. E. Amer. (U. S.)
**H. faunus** *Say.*—N. E. Amer.
**H. convivus** *Lec.*—N. J. Rare.
**H. vagans** *Lec.*—N. E. Amer.
**H. pennsylvanicus** *Dej.*—N. E. Amer. [U. S.]
*var.* **compar** *Lec.*—N. E. Amer.
*var.* **erythropus** *Dej.*—N. E. Amer.
**H. spadiceus** *Dej.*—N. J. [rare] and westward.
**H. fallax** *Lec.*—N. Y. [Linell], N. J.
**H. pleruiticus** *Kby.*—N. Y., N. J. and westward.
**H. herbivagus** *Say.*—N. E. Amer.
**H. nitidulus** *Chd.*—N. Y., N. J. Rare at seashore.
**H. innocuus** *Lec.*—Taken in this vicinity by Mr. Linell.
**H. rufimanus** *Lec.*—Pa.
**H. gravis** *Lec.*—Tex. [On the Buffalo list.]

## Selenophorus *Dej.*

**S. pedicularius** *Dej.*—N. Y. and southward. Seashore in Sept.
**S. iripennis** *Say.*—Ill., Ga., Tex. [On the Buffalo list.]
**S. gagatinus** *Dej.*—Mass. and westward.
**S. opalinus** *Lec.*—N. E. Amer. Seashore, Sept.
**S. ovalis** *Dej.*—N. J., Ga., Fla. Seashore, Sept.
**S. ellipticus** *Dej.*—N. Y., N. J., Ga., Tex. Seashore, Sept.

## Stenolophus *Dej.*

**S. carbonarius** *Brulle.*—Middle and Southern States. Seashore, Sept.
**S. spretus** *Dej.*—N. Y. and southward.
**S. fuliginosus** *Dej.*—N. Y., N. J. and westward.
**S. plebeius** *Dej.*—N. Y., N. J. Seashore, Sept.
**S. conjunctus** *Say.*—N. E. Amer. [U. S.]
**S. ochropezus** *Say.*—N. E. Amer.
**S. dissimilis** *Dej.*—Can., N, J., southward and westward.
**S. alternans** *Lec.*—Pa.

### ACUPALPUS *Lat.*

**A. hydropicus** *Lec.*—N. Y., N. J. May.
**A. carus** *Lec.*—Can., N. J., N. Y.

### BRADYCELLUS *Er.*

**B. linearis** *Lec.*—Pa., Wisc.
**B. cognatus** *Gyll.*—Europe and Northern America.
**B. neglectus** *Lec.*—Buffalo and northwest.
**B. rupestris** *Say.*—N. E. Amer.
**B. tantillus** *Dej.*—U. S.
**B. nigriceps** *Lec.*—N. Y., Va.

### TACHYCELLUS *Moraw.*

**T. nigrinus** *Dej.*—Alaska to Can.
**T. Kirbyi** *Horn.*—Canada to Ohio.
**T. atrimedius** *Say.*—N. E. Amer.
**T. badiipennis** *Hald.*—N. E. Amer.

### ANISODACTYLUS *Dej.*

**A. dulcicollis** *Laf.*—N. J. and southward.
**A. rusticus** *Say.*—N. E. Amer. Common everywhere.
**A. carbonarius** *Say.*—N. E. Amer.
**A. interpunctatus** *Kby.*—N. E. Amer.
**A. agricola** *Say.*—N. Y., southward and westward.
**A. Harrisii** *Lec.*—N. E. Amer.
**A. nigerrimus** *Dej.*—N. E. Amer.
**A. nigrita** *Dej.*—N. E. Amer.
**A. melanopus** *Hald.*—N. J., Pa., Ill.
**A. discoideus** *Dej.*—N. E. Amer. Rare.
**A. baltimorensis** *Say*,—N. E. Amer. Common.
**A. laetus** *Dej.*—N. Y., N, J., Ga., Tex.
**A. coenus** *Say.*—N. E. Amer.
**A. sericeus** *Harr.*—N. E. Amer.
**A. lugubris** *Dej.*—N. J.. Ia., Can., Mo.
**A. interstitialis** *Say.*—N. E. Amer.
**A. verticalis** *Lec.*—N. E. Amer.
**A. terminatus** *Say.*—N. Y., N. J. and westward.
**A. piceus** *Lec.*—N. E. Amer.
**A. semipunctatus** *Lec.*—Or. [On the Buffalo list.]
**A. punctulatus** *Lec.*—Middle State Region.

(TO BE CONTINUED.)